

Urgent treatment of spontaneous carotid artery pseudoaneurysm with covered stent graft: A case report

Spontan karotis arter psödoanevrizmasının kaplı stent ile acil tedavisi: Olgu sunumu

Engin Akgül , Gülen Sezer Alptekin Erkul , Sinan Erkul , Ahmet Hakan Vural 

Department of Cardiovascular Surgery, Dumlupınar University Evliya Çelebi Training and Research Hospital, Kütahya, Turkey

ABSTRACT

Pseudoaneurysms usually occur due to blunt trauma to the atherosclerotic vessels. The main pathophysiology is the disruption of the vessel wall and subsequent collection of the blood under adventitia and surrounding tissues. The localization of the lesion is determinant for the severity of the clinical presentation. Herein, we present a 64-year-old male case in whom Doppler ultrasound revealed a carotid artery pseudoaneurysm treated with endovascular method using a covered stent graft.

Keywords: Carotid artery pseudoaneurysm; covered stent graft; endovascular treatment.

ÖZ

Psödoanevrizmalar genellikle aterosklerotik damarlarda künt travmaya bağlı olarak ortaya çıkar. Ana patofizyoloji, damar duvarının bozulması ve sonrasında adventisya ve çevre dokular altında kanın toplanmasıdır. Lezyonun yerleşim yeri klinik tablonun şiddetini belirler. Bu yazıda, Doppler ultrasonda karotis arterde psödoanevrizma tespit edilen ve kaplı stent greft kullanılarak endovasküler yöntemle tedavi olarak tedavi edilen 64 yaşında bir erkek olgu sunuldu.

Anahtar sözcükler: Karotis arter psödoanevrizması; kaplı stent; endovasküler tedavi.

After the arterial wall injury which causes extravasation of the blood the damaged arterial wall is covered with a pseudocapsule defined as a pseudoaneurysm. Following the arterial wall disruption, hematoma or pseudoaneurysm formation occurs due to the arterial pressure. Although carotid pseudoaneurysms are rarely seen, they can cause catastrophic complications. In addition to the hemorrhagic and neurological effects, serious dyspnea may occur due to the mass effect of the pseudoaneurysm, such as the deviation of the trachea, and urgent intubation is required in such cases.

In this article, we present a case with a spontaneous left common carotid artery pseudoaneurysm with an

expanding hematoma treated with an endovascular stent graft in the emergency setting.

CASE REPORT

A 64-year-old male patient was admitted to the Evliya Çelebi Training and Research Hospital with dyspnea and neck pain for two days. On his physical examination, there was edema, hyperemia, and a pulsatile mass on the left side of the neck. Respiratory rate was 18 per min, the arterial blood pressure was 220/130 mmHg, and the heart rate was 110 bpm. The Glasgow Coma Scale score was 12 (eye: 3, verbal: 4, motor: 5). There was a left carotid artery pseudoaneurysm detected with color Doppler

Received: March 09, 2018 **Accepted:** March 27, 2018

Correspondence: Engin Akgül, MD, Dumlupınar Üniversitesi Evliya Çelebi Eğitim ve Araştırma Hastanesi, Kalp ve Damar Cerrahisi Kliniği, 43040 Kütahya, Turkey.
e-mail: engin_akgul@hotmail.com

Citation:

Akgül E, Alptekin Erkul GS, Erkul S, Vural AH. Urgent treatment of spontaneous carotid artery pseudoaneurysm with covered stent graft: A case report. Damar Cer Derg 2019;28(x):i-iii.



Figure 1. Angiographic view of pseudoaneurysm.



Figure 2. Treatment of pseudoaneurysm with covered stent graft.

ultrasonography. The patient was hospitalized and antihypertensive therapy was arranged. Right-sided central venous catheterization was applied. Although surgery was the first-line treatment of choice, clamping was not appropriate, as the mass grew rapidly in hours. Therefore, the patient was prepared for endovascular treatment and transferred to the angiography laboratory. The right femoral artery was cannulated with a 7-French (F) sheath under local anesthesia and 5000 IU of heparin injected by this way. The common carotid artery was cannulated with a 0.18 guidewire and a 6-F right coronary catheter. An 8x7-cm pseudoaneurysm was diagnosed at the pre-bifurcation of the internal and external carotid artery (Figure 1). An 8x59-mm covered stent graft (Atrium Advanta balloon expandable covered stent, USA) was implanted from the common carotid artery to the left internal carotid artery (Figure 2). The external carotid artery was also occluded with the covered stent graft due to the lesion localization. The patient was followed in the intensive care unit for one day and he was discharged on Day 4 without any neurological or vascular complications with daily acetylsalicylic acid 100 mg and clopidogrel 75 mg.

DISCUSSION

Carotid artery pseudoaneurysms are very rarely seen. The major causes are atherosclerosis,^[1] congenital

connective tissue disease, and traumas.^[2] In certain cases, carotid artery endarterectomy operation may also cause pseudoaneurysm formation^[2,3] as a result of suture line corruption.^[1] Arteritis, infection, arterial intimal degeneration, irradiation are the other causes.^[1,4] Carotid artery pseudoaneurysms are mostly located between the internal carotid artery and the common carotid artery,^[4] although there is a limited number of data about why it is most often seen at this level. As there is an increased risk of rupture and neurological complications for carotid artery pseudoaneurysms, they should be treated immediately.^[4]

The treatment modalities are open surgery and endovascular therapy. Primary repair of the pseudoaneurysm or using a patch for the repair are the open surgical techniques. Open surgery is the traditional treatment modality^[5] and is curative; however, exploring the pseudoaneurysm and clamping the responsible artery may be sometimes challenging. In particular, the risk of hemorrhage cannot be ruled out.^[5] Also, the cranial nerve injuries, usually of the IX, X, XII nerves, are some of the other serious complications which may occur during open surgery.^[6]

Recent studies demonstrate that endovascular treatment is less invasive and can be life-saving.^[5,7] For the complex lesions, endovascular treatment provides a relatively low risk and is as effective as open surgery.^[8] A systematic review of 113 studies with 224 patients

reported the midterm results for stroke, cranial nerve injury, and mortality rates as 1.8%, 0.5%, and 4.1% respectively with endovascular treatment.^[6]

In conclusion, endovascular treatment is technically feasible, suitable and favorable with a high procedure success. As the present case was admitted with giant neck swelling and short neck, open surgery would be difficult and we preferred endovascular treatment. We believe that we make the right choice based on the health status of the patient and in the light of current literature data.

Declaration of conflicting interests

The authors declared no conflicts of interest with respect to the authorship and/or publication of this article.

Funding

The authors received no financial support for the research and/or authorship of this article.

REFERENCES

1. Ayoub K, Nacion KM, Bartholomew JR. Carotid artery pseudoaneurysm. *JVU* 2010;34:195-8.
2. Fankhauser GT, Stone WM, Fowl RJ, O'Donnell ME, Bower TC, Meyer FB, et al. Surgical and medical management of extracranial carotid artery aneurysms. *J Vasc Surg* 2015;61:389-93.
3. Srivastava SD, Eagleton MJ, O'Hara P, Kashyap VS, Sarac T, Clair D. Surgical repair of carotid artery aneurysms: a 10-year, single-center experience. *Ann Vasc Surg* 2010;24:100-5.
4. Garg K, Rockman CB, Lee V, Maldonado TS, Jacobowitz GR, Adelman MA, et al. Presentation and management of carotid artery aneurysms and pseudoaneurysms. *J Vasc Surg* 2012;55:1618-22.
5. Tsurukiri J, Okumura E, Yamanaka H, Jimbo H, Hoshiai A. Emergency endovascular treatment of a ruptured external carotid artery pseudoaneurysm caused by a cervical stab wound: A case report and literature review. *Ulus Travma Acil Cerrahi Derg* 2017;23:433-7.
6. Garg K, Rockman CB, Lee V, Maldonado TS, Jacobowitz GR, Adelman MA, et al. Presentation and management of carotid artery aneurysms and pseudoaneurysms. *J Vasc Surg* 2012;55:1618-22.
7. Li Z, Chang G, Yao C, Guo L, Liu Y, Wang M, et al. Endovascular stenting of extracranial carotid artery aneurysm: a systematic review. *Eur J Vasc Endovasc Surg* 2011;42:419-26.
8. Kadyrov NA, Friedman JA, Nichols DA, Cohen-Gadol AA, Link MJ, Piepgras DG. Endovascular treatment of an internal carotid artery pseudoaneurysm following transsphenoidal surgery. Case report. *J Neurosurg* 2002;96:624-7.